PTO/SB/08A(10-01)
Approved for use through 10/31/2002, OMB 631-0031
US Patent & Trademark Office: U.S. DEPARTHEENT OF COMMERCE
n of information unless & contains a valid OMB control number.

Substitute for form 1449A/PTO Complete if Known INFORMATION DISCLOSURE 09/765,063 **Application Number** STATEMENT BY ADMINISTRATE (Use as many sheets as necessary) **Filing Date** January 17, 2001 Yaksh, Tony **First Named Inventor** Unknown **Group Art Unit** Roy, Baisakhi **Examiner Name** Attorney Docket No: 1133.029US1 Sheet 1 of 4

	OTHER	R DOCUMENTS - NON PATENT LITERATURE DOCUMENTS	
Examiner Initials*	Cite No 1	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	T²
		ABBADIE, C., et al., "Differential Contribution of the Two Phases of the Formalin	
8·R·	 	Test to the Pattern of c-fos Expression in the Rat Spinal Code: Studies With	
ייע		Remifentanil and Lidocaine", Pain, 69, (1997),101-110	
1		ABBOTT, F. V., et al., "The Formalin Test: Scoring Properties of the First and	
Ì		Second Phases of the Pain Response in Rats", Pain, 60, (1995),91-102	
		ABRAM, S. E., et al., "Morphine, but not Inhalation Anaesthesia, Blocks Post-	
1		Injury Facilitation. The Role of Preemptive Suppression of Afferent	
- 1		Transmission", Anesthesiology, 78(4), (1993),713-721	
		ABRAM, S. E., et al., "Systemic Lidocaine Blocks Nerve Injury-Induced	
1		Hyperalgesia and Nociceptor-Driven Spinal Sensitization in the Rat",	
1		Anesthesiology, 80(2), (1994),383-391	
		ALOISI, A. M., et al., "Behavioral Effects of Different Intensities of Formalin Pain	
1		in Rats", Physiology & Behavior, 58(3), (1995),603-610	
		BANNON, A. W., et al., "ABT-594 [(R)-5-(2-azetidinylmethoxy)-2-chloropyridine]:	
1	.	A Novel, Orally Effective Antinociceptive Agent Acting via Neuronal Nictotinic	
!		Acetylcholine Receptors: II. In Vivo Characterization", The Journal of	
		Pharmacology and Experimental Therapeutics, 285(2), (1998),787-794	
-		BHATNAGAR, S., et al., "The Effects of Prior Chronic Stress on Cardiovascular	
		Responses to Acute Restraint and Formalin Injection", Brain Research, 797,	
1		(1998),313-320	
		BRENNAN, T. J., "AMPA/Kainate Receptor Antagonists as Novel Analgesic	
		Agents", Anesthesiology, 89(5), (1998),1049-1051	
		BUERKLE, H., et al., "Effect of Continuous Spinal Remifentanil Infusion on	
-		Behaviour and Spinal Glutamate Release Evoked by Subcutaneous Formalin in	
		the Rat", British Journal of Anaesthesia, 80, (1998),348-353	
		CHAPLAN, S. R., et al., "Efficacy of Spinal NMDA Receptor Antagonism in	1
1		Formalin Hyperalgesia and Nerve Injury Evoked Allodynia in the Rat", <u>Journal of</u>	
i i		Pharmacology and Experimental Therapeutics, 280(2), (1997),829-838	
 		CLAVELOU, P., et al., "The Orofacial Formalin Test in Rats: Effects of Different	
		Formalin Concentrations", Pain, 62, (1995),295-301	
		CODERRE, T. J., et al., "The Formalin Test: A Validation of the Weighted-	T^{-}
		Scores Method of Behavioral Pain Rating", Pain, 54, (1993),43-50	
		DALLEL, R., et al., "Evidence for a Peripheral Origin of the Tonic Nociceptive	T
		Response to Subcutanous Formalin", Pain, 61, (1995),11-16	
		DICKENSON, A. H., et al., "Peripheral Origins and Central Modulation of	1
		Subcutaneous Formalin-Induced Activity of Rat Dorsal Horn Neurones",	
		Neuroscience Letters, 83, (1987),207-211	
+		DICKERSON, A. H., et al., "Chapter 36 Response Properties of Dorsal Horn	
ı		Neurons: Pharmacology of the Dorsal Horn", In: Anesthesia: Biologic	
0 0		Foundations, Yaksh, T., et al., Editors, Lippincott-Raven Publishers, (1997),611-	
B·R·		624	1

EXAMINER	barsakhi	Loy	DATE CONSIDERED	6-20-05
----------	----------	-----	-----------------	---------

PTC/SB/084 (10-01)
Approved for use through 10/31/2002, ONIB 651-0031
US Prient & Tridement Office: U.S. DEPARTMENT OF COMMERCE

Substitute for form 1449A/PTO	Complete If Known	Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control numb. Complete If Known		
INFORMATION DISCLOSURE STATEMENT BY APPLICANT	Application Number	09/765,063		
(Use as many sheets as necessary)	Filing Date	January 17, 2001		
	First Named Inventor	Yaksh, Tony		
	Group Art Unit	Unknown		
Sheet 2 of 4	Examiner Name	Roy, Baisakhi		
	Attorney Docket No: 1	1133.029US1		

Examiner Initials*	Cite No 1	include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s),	T
		publisher, city and/or country where published.	L
		DIRIG, D. M., et al., "Intrathecal Baclofen and Muscimol, but not Midazolam, are	1
B. R.		Antinociceptive Using the Rat-Formalin Model", Journal of Pharmacology and	
		Experimental Therapeutics, 275(1), (1995),219-227	╙
1		DRAY, A., et al., "Systemic Capsaicin and Olvanil Reduce the Acute Algogenic	
1 1		and the Late Inflammatory Phase Following Formalin Injection into Rodent Paw",	
		Pain, 47, (1991),79-83	L
		DUBUISSON, D., et al., "The Formalin Test: A Quantitative Study of the	
		Analgesic Effects of Morphine, Meperidine, and Brain Stem Stimulation in Rats	
		and Cats", <u>Pain, 4,</u> (1977),161-174	<u> </u>
		HANDWERKER, H. O., "Electrophysiological Mechanisms in Inflammatory	
		Pain", Agents and Actions. Supplements, 32 - Drugs in Inflammation, (1991),91-	
		99	L
		HUNTER, J. C., et al., "Role of Excitatory Amino Acid Receptors in the Mediation	
- 1 1		of the Nociceptive Response to Formalin in the Rat", Neurosci Lett, 174,	
		(1994),217-221	L
		JETT, M. F., et al., "The Formalin Test in Rat: Validation of an Automated	
		System", Pain, 64, (1996),19-25	<u> </u>
	•	JOURDAN, D., et al., "A New Automated Method of Pain Scoring in the	
		Formalin Test in Rats", <u>Pain, 71,</u> (1997),267-270	┡
		MALMBERG, A. B., et al., "Antinociceptive Actions of Spinal Nonsteroidal Anti-	ł
		Inflammatory Agents on the Formalin Test in the Rat", The Journal of	
		Pharmacology and Experimental Therapeutics, 263(1), (1992),136-146	┺
		MALMBERG, A. B., et al., "Antinociceptive Effect of Spinally Delivered	
		Prostaglandin E Receptor Antagonists in the Formalin Test on the Rat",	
		Neuroscience Letters, 173, (1994),193-196	├_
		MALMBERG, A. B., et al., "Effect of Continuous Intrathecal Infusion of ω -	
		conopeptides, N-type Calcium-Channel Blockers, on Behavior and	
1 1		Antinociception in the Formalin and Hot-Plate Tests in Rats", Pain, 60,	
		(1995),83-90	-
		MALMBERG, A. B., et al., "Pharmacology of the Spinal Action of Ketorolac,	
		Morphine, ST-91, U50588H, and L-PIA on the Formalin Test and an	
		Isobolographic Analysis of the NSAID Interaction", Anesthesiology, 79(2).	
		(1993),270-281	-
		MALMBERG, A. B., et al., "Spinal Nitric Oxide Synthesis Inhibition Blocks	
		NMDA-Induced Thermal Hyperalgesia and Produces Antinociception in the	
		Formalin Test in Rats", Pain, 54, (1993),291-300 NOZAKI-TAGUCHI, N., et al., "A Novel Model of Primary and Secondary	╁
		Hyperalgesia After Mild Thermal Injury in the Rat", Neuroscience Letters, 254.	
B.R.		(1998),25-28	
יאיטו		(1330),20-20	十

EXAMINER	baisaphi	Lay	DATE CONSIDERED	6-20-05
----------	----------	-----	-----------------	---------

PTO/SB/08A(10-01)
Approved for use through 10/31/2002, OMB 651-0031
US Patent's Tradement Global V.S. DEFARTMENT of COLUMENCE
Inequired to respond to a collection of information unless it contrals a valid OMB control number.

Examiner Name	Roy, Baisakhi		
Group Art Unit	Unknown		
First Named Inventor	Yaksh, Tony		
Filing Date	January 17, 2001		
Application Number	09/765,063		
Complete if Known			
	Application Number Filing Date First Named Inventor Group Art Unit		

Examiner No.* Include name of the author (in CAPITAL LETTERS), tille of the article (when appropriate), tille of the liem (brook, magatine, burnal, serial, sympostum, catalog, etc.), date, pagele), volume-lessue number(s), published. PETERSON, M. A., et al., "The Differential Contribution of Capsaicin-Sensitive Afferents to Behavioral and Cardiovascular Measures of Brief and Persistent Nociception and to Fos Expression in the Formalin Test", Brain Research, 755(1), (1997),9-16 PRADO, W. A., et al., "Antinociceptive Effect of Intrathecal Neostigmine Evaluated in Rats by Two Different Pain Models", Brazilian Journal of Medical and Biological Research, 30, (1887),1225-1231 PRICE, D. D., "Psychophysical Observations on Patients With Neuropathic Pain Relivered by a Sympathetic Block", Pain, 36, (1989),273-288 PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Ffects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Keinate GluRS Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neurophymmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., "Pany Pany Pany Pany Pany Pany Pany Pany		OTHER	R DOCUMENTS - NON PATENT LITERATURE DOCUMENTS	
Afferents to Behavioral and Cardiovascular Measures of Brief and Persistent Nociception and to Fos Expression in the Formalin Test", Brain Research, 755(1), (1997),9-16 PRADO, W. A., et al., "Antinociceptive Effect of Intrathecal Neostigmine Evaluated in Rats by Two Different Pain Models", Brazilian Journal of Medical and Biological Research, 30, (1887),1225-1231 PRICE, D. D., "Psychophysical Observations on Patients With Neuropathic Pain Relivered by a Sympathetic Block", Pain, 36, (1989),273-288 PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., "Fairly Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanii", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),757-57-584 TAYLOR, B. K., et al., "Privilary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "T			(book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	T
Nociception and to Fos Expression in the Formalin Test*, Brain Research, 755(1), (1997),9-16 PRADO, W. A., et al., "Antinociceptive Effect of Intrathecal Neostigmine Evaluated in Rats by Two Different Pain Models*, Brazilian Journal of Medical and Biological Research, 30, (1887),1225-1231 PRICE, D. D., "Psychophysical Observations on Patients With Neuropathic Pain Relivered by a Sympathetic Block*, Pain, 36, (1989),273-288 PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity*, Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat*, Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test*, Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate Gluns Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat*, Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine*, Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Presistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Privitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology,			PETERSON, M. A., et al., "The Differential Contribution of Capsaicin-Sensitive	
755(1), (1997),9-16 PRADO, W. A., et al., "Antinociceptive Effect of Intrathecal Neostigmine Evaluated in Rats by Two Different Pain Models", Brazilian Journal of Medical and Biological Research, 30, (1887),1225-1231 PRICE, D. D., "Psychophysical Observations on Patients With Neuropathic Pain Relivered by a Sympathetic Block", Pain, 36, (1989),273-288 PUIG, S., et al., "Formalin-Evoked Activity in Identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),757-5784 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of t	0.0		Afferents to Behavioral and Cardiovascular Measures of Brief and Persistent	
PRADO, W. A., et al., "Antinociceptive Effect of Intrathecal Neostigmine Evaluated in Rats by Two Different Pain Models", Brazilian Journal of Medical and Biological Research, 30, (1887),1225-1231 PRICE, D. D., "Psychophysical Observations on Patients With Neuropathic Pain Relivered by a Sympathetic Block", Pain, 36, (1989),273-288 PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate Gluf5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanii", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJÄLSEN, A., et al., "The Formalin Test: An Evaluation of the Metho	β·κ·		Nociception and to Fos Expression in the Formalin Test", <u>Brain Research</u> ,	ł
Evaluated in Rats by Two Different Pain Models", Brazilian Journal of Medical and Biological Research, 30, (1887),1225-1231 PRICE, D. D., "Psychophysical Observations on Patients With Neuropathic Pain Relivered by a Sympathetic Block", Pain, 36, (1989),273-288 PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Presistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "The Rat Paw Formal	i i			<u> </u>
and Biological Research, 30, (1887),1225-1231 PRICE, D. D., "Psychophysical Observations on Patients With Neuropathic Pain Relivered by a Sympathetic Block", Pain, 36, (1989),273-288 PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanii", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Pharmacology, 13(1998),2407-2413 TJØLSEN, A., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			PRADO, W. A., et al., "Antinociceptive Effect of Intrathecal Neostigmine	
PRICE, D. D., "Psychophysical Observations on Patients With Neuropathic Pain Relivered by a Sympathetic Block", Pain, 36, (1989),273-288 PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),757-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of	i			1
Relivered by a Sympathetic Block", Pain, 36, (1989),273-288 PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			and Biological Research, 30, (1887),1225-1231	
PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers: Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Presistent Cardiovascular Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test". An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44				
Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355 RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat. Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of	Į.		Relivered by a Sympathetic Block", Pain, 36, (1989),273-288	
RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			PUIG, S., et al., "Formalin-Evoked Activity in identified Primary Afferent Fibers:	
Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of	1		Systemic Lidocaine Suppresses Phase-2 Activity", Pain, 64, (1995),345-355	<u> </u>
Trigeminal Brain Stem Nociceptive Neurones in the Rat", Journal of Neurophysiology, 73(2), (1995),496-505 SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			RABOISSON, P., et al., "Effects of Subcutaneous Formalin on the Activity of	
SHIMOYAMA, N., et al., "Spinal Gabapentin is Antinociceptive in the Rat Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44			Trigeminal Brain Stem Nociceptive Neurones in the Rat", <u>Journal of</u>	
Formalin Test", Neuroscience Letters, 222(1), (1997),65-67 SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanii", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44			Neurophysiology, 73(2), (1995),496-505	<u> </u>
SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of				
Nociceptive Response to Formalin in the Rat", Neuropharmacology, 37, (1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			Formalin Test", Neuroscience Letters, 222(1), (1997),65-67	ļ
(1998),25-36 SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			SIMMONS, R. M., et al., "Kainate GluR5 Receptor Subtype Mediates the	
SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of	1			
Anxiolytic-Like and Antinociceptive Actions that are Reversed by D-Serine", Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of	}		(1998),25-36	ļ <u>.</u>
Psychopharmacology, 127, (1996),1-9 TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			SINGH, L., et al., "The Antiepileptic Agent Gabapentin (Neurontin) Possesses	1
TALLARIDA, R. J., et al., Manual of Pharmacologic Calculations With Computer Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of	l			
Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291 TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of				
TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile, but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of				
but not the Magnitude, of the Tonic Response to Subcutaneous Formalin: Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			Programs, (2nd Edition, 1987, Springer-Verlag New York, Inc.),Pg. 291	
Effects with Remifentanil", The Journal of Pharmacology and Experimental Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			TAYLOR, B. K., et al., "Early Nociceptive Events Influence the Temporal Profile,	
Therapeutics, (1997),876-883 TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			but not the Magnitude, of the Tonic Response to Subcutaneous Formalin:	
TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of				
Responses to Subcutaneous Formalin Require Peripheral Nerve Input", The Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			<u>Therapeutics</u> , (1997),876-883	ļ
Journal of Neuroscience, 15(11), (1995),7575-7584 TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			TAYLOR, B. K., et al., "Persistent Cardiovascular and Behavioral Nociceptive	
TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of	1			
Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			Journal of Neuroscience, 15(11), (1995),7575-7584	<u> </u>
Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413 TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			TAYLOR, B. K., et al., "Pituitary-Adrenocortical Responses to Persistent Noxious	1
TJØLSEN, A., et al., "The Formalin Test: An Evaluation of the Method", Pain, 51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			Stimuli in the Awake Rat: Endogenous Corticosterone Does Not Reduce	
51, (1992),5-17 WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			Nociception in the Formalin Test", Endocrinology, 139(5), (1998),2407-2413	<u> </u>
WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of				
for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44 WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			51, (1992),5-17	₩
WHEELER-ACETO, H., et al., "The Rat Paw Formalin Test: Comparison of			WHEELER-ACETO, H., et al., "Standardardization of the Rat Paw Formalin Test	1
			for the Evaluation of Analgesics", Psychopharmacology, 104, (1991),35-44	ļ
Noxious Agents", <u>Pain, 40,</u> (1990),229-238	() n			
i i	P.K.		Noxious Agents", <u>Pain, 40, (1990),229-238</u>	<u> </u>
				<u> </u>

EXAMINER	bousakhi	Lan	DATE CONSIDERED	6-20-05
	UXXXXXXXXXXXXXXX	'ALLIA		• • • •

PTO/SB/08A(10-01)
Approved for use through 10/31/2002, OMB 651-0031

Substitute for form 1449A/PTO	Complete & Known			
INFORMATION DISCLOSURE STATEMENT BY APPLICANT	Application Number	09/765,063		
(Use as many sheets as necessary)	Filing Date	January 17, 2001		
	First Named Inventor	Yaksh, Tony		
	Group Art Unit	Unknown		
	Examiner Name	Roy, Baisakhi		
Sheet 4 of 4	Attorney Docket No: 1	133.029US1		

	OTHE	R DOCUMENTS NON PATENT LITERATURE DOCUMENTS	
Examiner initials*	Cite No ¹	include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	T
B · R ·		WOOLF, C. J., et al., "Long Term Alterations in the Excitability of the Flexion Reflex Produced by Peripheral Tissue Injury in the Chronic Decerebrate Rat", Pain, 18, (1984),325-343	
		YAKSH, T. L., et al., "Central Pharmacology of Nociceptive Transmissions", In: Textbook of Pain, (4th Edition, Churchill Livingstone),(1999),253-308	
		YAKSH, T. L., "Chapter 40 Preclinical Models of Nociception", In: Anesthesia: Biologic Foundations (Vol. 1), Yaksh, T., et al., Editors, Lippincott-Raven Publishers, (1997), 685-718	
		YAKSH, T. L., "Chronic Catherization of the Spinal Subarachnoid Space", Physiology & Behavior, 17, (1976),1031-1036	
		YAKSH, T. L., et al., "Intrathecal Capsaicin Depletes Substance P in the Rat Spinal Cord and Produces Prolonged Thermal Analgesia", Science, 206(4417), (1979),481-483	
		YAKSH, T. L., et al., "The Spinal Biology in Humans and Animals of Pain States Generated by Persistent Small Afferent Input", Pro. Natl. Acad. Sci. USA, 96, (1999),7680-7686	
	-	YAMAMOTO, T., et al., "Stereospecific Effects of a Nonpeptidic NK1 Selective Antagonist, CP-96,345: Antinociception in the Absence of Motor Dysfunction", Life Sciences, 49(26), (1991),1955-1963	
		YOON, M. H., et al., "The Effect of Intrathecal Gabapentin on Pain Behavior and Hemodynamics on the Formalin Test in the Rat", <u>Anesthesia and Analgesia</u> , 89, (1999),434-439	
B. R.		ZAR, J. H., "Table of Contents", <u>Biostatistical Analysis</u> , (2nd Edition, 1984, Prentice Hall, Inc.),134-138	

EXAMINER barsakhi Loy DATE CONSIDERED 6-20-05